

## Interim Grass fed milk Standard Protocol

The temporary grass-fed standard has developed by DFO in order to take advantage of the 2017 grazing season. This temporary standard was being developed for Ontario but the goal is to work with provinces to have a national standard conjunction with provinces.

This temporary standard is open for consultation and the next steps are as following:

1-From June 2017 to March 2019: The temporary standard is in place.

2- From June 2017 -August 2018:

- Consultation period: Please send your comments and feedback to Bita Farhang  
Bita.Farhang@milk.org

3- From November 2017 -October 2018:

- Review all comments and feedback
- Conduct a Bio markers level analysis

4- National Standard submission to CFIA: March 2019

The DFO standard is being referred to as a “temporary” standard as the Canadian standard will be used exclusively when reviewed by the CFIA.

Note: Since DFO started the development of a grass-fed standard, a number of other provinces also expressed interest in a grass-fed standard and as a result of this interest Dairy Farmers of Canada agreed to lead the development of a Canadian grass-fed standard using DFO’s grass-fed standard as a template with a goal to have a national standard in place, including CFIA review, for the 2018 grazing season. DFC established a small technical working group in December 2017. The group is made up of provincial members, DPAC, technical experts etc. to proceed with developing a final standard for submission to CFIA. The technical committee in June 2018 decided to conduct 2 studies: one on consumer perception, and the other on the cost of producing grass-fed milk. Due to the current differing methods of grass-fed production, the studies will be important in determining which method is chosen as part of the final standard.

In the case of consumer perception, the goal of the study is to understand what consumers expect when reading the “grass-fed” label – particularly when it comes to animal welfare or access to pasture. Depending on how important each attribute is based on the study will determine how the standards will be edited.

The consumer study will take a few months to complete, and the report is projected for some time in early fall. As for the COP study, the final report is expected in December. In the meantime, DFC will work on exploring the logo requirements. Once received, the committee can then review the results and edit the standard in early 2019, leaving some room for a final consultation and submission to CFIA targeted for March 1st, 2019.

# Dairy Farmers of Ontario's Interim Grass-Fed Milk Standard

During the last couple of years, there has been an increasing number of demands from processors for the supply of grass-fed milk. At this time, there are no established national or provincial standards for grass-fed milk in Canada. This document aims at setting the basis of the Canadian standards for Grass-fed milk.

Efficient grass-fed farming requires a high forage diet with as much fresh grass as possible with limited amount of concentrates, considering limitations imposed by climate and other environmental factors and cow's nutritional needs. Cows in early lactation may need more concentrates than cows in mid lactation and cows in late lactation may not need to be fed any concentrates. For this reason, managing a grass-fed operation requires commitment to the operation and to grass-fed principles.

This Protocol focusses primarily on feed and farming requirements. A secondary focus is to authenticate grass fed milk based on specific biomarkers that are metabolized by cows under a grass-fed diet. Proposed biomarkers are a minimum level of conjugated linoleic acid (CLA) and a maximum ratio of Omega 6: Omega 3 fatty acids. It is important to note that the biomarkers testing is considered as a management tool for monitoring the feeding requirements instead of absolute values to reach or not to exceed, regardless of the type of feeds used.

## 1. Feeding Protocol

- 1.1. Forage/grass must provide at least 75 % of the total dry matter intake of a dairy cow.
- 1.2. 25% of the total dry matter intake is targeted to come from grains and supplements except prohibited feeds, as listed in 1.3. This ratio can go up to maximum 30% of the total dry matter intake in non-growing season with a written recommendation from a nutritionist.
- 1.3. Grass-fed animals **cannot be fed the prohibited feeds** listed below:
  - **Corn silage**, but can be fed corn grain
  - **Corn distiller grains**
  - **Any type of plant oils/fats**
  - **Any type of Marine oils/fats/by-products**
  - **Any type of animal or poultry fat**
  - **Full-fat soybeans, sunflower and safflower seeds or oils but can be fed these meals**
  - **Linseed (flaxseed) for less than 1 kg per day.**
  - **Urea or any other non-protein nitrogen supplements.**
- 1.4. Mineral and vitamin supplements may be provided as required (as prescribed by the manufacturer, a veterinarian or qualified nutritionist) with the exception of mineral and vitamin supplements containing any prohibited ingredients (e.g., fishmeal, and urea).
- 1.5. A record of feeding and pasture protocol must be kept on a weekly basis, and the grass-fed farm must fully keep all documents
- 1.6. Forage is defined as "any herbaceous plant material that can be grazed or harvested for feeding, except for post-vegetative state cereal/grain". This includes grass (annual and perennial), forbs (legume and brassicas), browse, and cereal grain crops in the vegetative state.

The vegetative state is a stage in cereal plant growth at which the plant “contains non-reproductive plant parts (leaf and stem), in contrast to reproductive plant parts (flower and seed) and is still in the developmental stages of plant growth.” According to Zadok’s table, which illustrates cereal grain growth and development, large grains, such as corn, should be in the development stage before the boot stage to ensure a vegetative state.

## 2. Pasture Management, Grazing, Confinement and Stock Piled Forages

- 2.1. All dairy cows should have access to pasture at least 120 days a year for 6 hours per day during the grazing period. In regions where the grazing period is less than 120 days, they must be on pasture for at least 6 hours per day while the weather permits.
- 2.2. A pasture management plan must be in place to show pasture is being managed as a crop and is being rotated to allow for proper rest and regrowth of the pasture before the animals are reintroduced to the area.
- 2.3. Pasture shall be managed in a way that will not degrade the soil or quality of the pasture and natural resources with the exception of having a sacrifice area during the non-grazing season.
- 2.4. The grazing area must be greater than 0.13 hectare (0.33 acre) per cow.

## 3. Animal Health and Welfare

The welfare of the animal always comes first. This Protocol requires compliance with the Canadian Code of Practice for the Care and Handling of Dairy Cattle. To be on the grass fed program, compliance with this Code must be validated through the proAction® program.

## 4. Auditing and verification

4.1 The feeding protocol and the biomarkers are verified by DFO during the temporary phase, but the grass-fed farms will be audited by a qualified and independent third party when the national standard is in place, to ensure compliance to the Protocol. This audit would cover:

- On-farm production conditions, which must be checked at least once a year
- The authentication of milk that is produced under this Protocol

4.2 Authentication of milk that is produced under this Protocol is based on the feeding protocol and specific biomarkers that are transferred from the diet to the milk or that are specifically produced by ruminal microorganisms or by the animals’ metabolism under a grass-fed diet as follows:

- The CLA in grass-fed milk is greater than or equal to 4 (mg-g-1 total fat).
- The 18:2 n-6/18:3 n-3 ratio in grass-fed milk is less than or equal to 2.5 (*it might change to less than or equal to 3 - this should be verified*)
- Note: The biomarker levels will be validated and can be modified accordingly. During the validation period, other biomarkers, such as the ratio of palmitic acid (C16) to oleic

acid (C18:1 cis-9) and the ratio of (C18:1 trans-11) to (C18:1 trans-10), will be also monitored.

- 4.3 Prior to supplying milk into a grass-fed program, the grass-fed farm must demonstrate that the biomarkers are tested in the milk on regulatory bulk tank samples weekly during the first four or 10 weeks of being on the grass-fed program, and at least monthly thereafter.
- 4.4 Milk cannot be marketed as grass-fed until two (2) consecutive bulk tank tests meet the minimum level of the biomarkers and a third-party auditor provides a report indicating compliance with the Protocol.

## 5. Definitions

**Boot Stage:** The flag leaf is fully expanded, but the awns and grain head are not visible. The grain head can be felt in the flag leaf sheath.

**Brassicas:** A family of very productive annual forage vegetables used as transition crops between pasture renovations or as a supplemental feed source for extending the grazing season when other forages are less productive. Examples include turnips, rape, and kale.

**Browse:** 1) Leaf and twig growth of shrubs, woody vines, trees, cacti, and other non-herbaceous vegetation available for animal consumption. 2) To browse: the consumption of browse in situ by animals (Barnes R.F. et al 1995).

**Diet:** The feed regularly offered to or consumed by an animal (Barnes R.F. et al 1995).

**Ensiled:** Having been subjected to anaerobic fermentation to form silage. (Cullison, A. E. 1979)

**Fermentation:** Chemical changes brought about by enzymes produced by various microorganisms. (Cullison, A. E. 1979)

**Forage:** Edible parts of plants, other than separated grain, that can provide feed for grazing animals, or that can be harvested for feeding. Includes browse, herbage, and mast; Vegetative material in a fresh, dried, or ensiled state (pasture, hay, or silage), which is fed to livestock.

**Forb:** Any herbaceous broadleaf plant that is not a grass and is not grass-like. (Barnes R.F. et al 1995)

**Fruit:** 1) The usually edible reproductive body of a seed plant, one having a sweet pulp associated with the seed. 2) A product of fertilization in a plant with its modified envelopes or appendages, specifically the ripened ovary of a seed plant and its contents.

**Grain:** Seed from cereal plants, caryopsis. Corn, wheat, rye, oats, rice, millet, sorghum, barley, triticale.

**Grass:** Member of the plant family Poaceae (Barnes R.F. et al 1995).

**Graze:** 1) the consumption of standing or residual forage by livestock; 2) to put livestock to feed on standing residual forage

**Grazing Season:** The period when pasture is available for grazing, due to natural precipitation or irrigation.

**Growing Season:** The number of days between the last spring freeze date and the first fall freeze date.

**Hay:** The aerial parts of forage crops stored in the dry form for animal feeding (Cullison A.E. 1979).

**Haylage:** Haylage is the feed produced by storing a forage crop; dried to a moisture level of about 45- 55% in an airtight silo (Cullison A.E. 1979).

**Legumes:** Members of the Fabaceae plant family (formerly known as the Leguminosae family). Legumes are dicots (produce two seed leaves), produce seed in a pod, have netted leaf venation, and usually have a taproot type of root system. Most legumes can interact with bacteria of the genus *Rhizobium* to fix

nitrogen in nodules on their roots. Legumes may have one of four types of seedheads. These seedhead types are the raceme, the spike, the head or umbel (Ball D.M. et al. 2007).

**Meadow:** Area covered with grasses and/or legumes, often native to the area, grown primarily for hay but with secondary grazing potential (Barnes R.F. et al 1995).

**Mineral:** 1) a solid homogeneous crystalline chemical element or compound that results from the inorganic processes of nature. 2) Any of the various naturally occurring homogeneous substances obtained usually from the ground. 3) a synthetic substance having the chemical composition and crystalline form and properties of a naturally occurring mineral.

**Non-Grazing Season:** The period when pasture is not available for grazing, due to natural precipitation or weather. Varies with the region of Canada.

**Pasture:** 1) Forages harvested by grazing animals. 2) An area of land with 75% forage cover or unbroken land on which livestock may graze at will.

**Sacrifice pasture:** A pasture where animals may move about and express their natural behaviours and where forage can be grown during the grazing season.

**Separated Grain:** Grain detached from cereal crop plants.

**Silage:** Forage preserved in a succulent condition by partial anaerobic, acid fermentation.

**Supplement:** A nutritional additive (salt, protein, phosphorus, etc.) intended to improve the nutritional balance and remedy deficiencies of the diet (Barnes R.F. et al 1995).

**Supplemental Feeding:** The practice of supplying feedstuffs to correct nutritional deficiencies in an animal's "natural" diet.


**Vegetative:** Non-reproductive plant parts, (leaf and stem) in contrast to reproductive plant parts (flower and seed) in developmental stages of plant growth. The non-reproductive stage in plant development (Barnes R.F. et al 1995).

**Vegetative State/Stage:** A stage in cereal plant growth at which the plant "contains non-reproductive plant parts (leaf and stem) in contrast to reproductive plant parts (flower and seed) and is still in the developmental stages of plant growth.

**Vitamin:** Any of various organic substances essential in minute quantities to the nutrition of most animals and some plants that act especially as coenzymes and precursors of coenzymes in regulating of metabolic processes

## APPENDIX A

Figure 1: Zadoks Scale, Which Describes the Stages of Grain Plant Growth and Development



## REFERENCES

1. American Grassfed Association. (2016). Grassfed&Grass Pastured Ruminant Standards.
2. Ball, D. M., Hoveland, C. S., & Lacefield, G. D. (2007). Southern Forages: Modern Concepts for Forage Crop Management 4th ed.
3. Barnes, R.F, Miller, D.A., Nelson, C. J. (1995). Forages, Vol. I, An Introduction to Grassland Agriculture: Glossary (5th ed.). Ames, IA: Iowa State University Press.
4. Benbrook, C. M., Butler, G., Latif, M. A., Leifert, C., & Davis, D. R. (2013). Organic production enhances milk nutritional quality by shifting fatty acid composition: A United States-wide, 18-month study. PLoS ONE, 8(12). <https://doi.org/10.1371/journal.pone.0082429>
5. Blasko J., Kubinec R., Gorova R., Fabry I., Lorenz W., Sojak L.(2010) Fatty acid composition of summer and winter cows' milk and butter. Journal of Food and Nutrition Research .49(4):169–177
6. Capuano, E., Van Der Veer, G., Boerrigter-Eenling, R., Elgersma, A., Rademaker, J., Sterian, A., & Van Ruth, S. M. (2014). Verification of fresh grass feeding, pasture grazing and organic farming by cows farm milk fatty acid profile. Food Chemistry, 164, 234–241. <https://doi.org/10.1016/j.foodchem.2014.05.011>
7. Chiliard, Y, Ferley A. Doreau , M.( 2001).Effect of different types of forages, animal fat or marine oils incow's diet on milk fat secretion and composition, especially conjugated linoleic acid (CLA) and polyunsaturated fatty acids Livestock Production Science 70 :31–48
8. Cullison, A. E. (1979). Feeds and Feeding.

9. Elgersma A., Wever A. C. and Nałęcz-Tarwacka T. (2006). Grazing versus indoor feeding: effects on milk quality, Sustainable Grassland Productivity: Proceedings of the 21st General Meeting of the European Grassland Federation Badajoz, Spain 3-6 April 2006 419-427. <http://edepot.wur.nl/571>
10. Ferly A., Glasser F., Martin B., Andueza D., Chilliard Y. (2011). Effects of Feeding Factors and Breed on Cow Milk Fatty Acid Composition: Recent Data Bulletin UASVM, Veterinary Medicine 68(1): 137-145.
11. R. Mohammed , C. S. Stanton , J. J. Kennelly , J. K. G. Kramer , J. F. Mee , D. R. Glimm , M. O'Donovan and J. J. Murphy(2009). Grazing cows are more efficient than zero-grazed and grass silage-fed cows in milk rumenic acid production. Journal of Dairy Science .92 ( 8),3874-3893.
12. O'Callaghan, T. F., Hennessy, D., McAuliffe, S., Kilcawley, K. N., O'Donovan, M., Dillon, P., Stanton, C. (2016). Effect of pasture versus indoor feeding systems on raw milk composition and quality over an entire lactation. Journal of Dairy Science, 99(12), 9424–9440. <https://doi.org/10.3168/jds.2016-10985>
13. Pennsylvania Certified Organic. (2000). Pennsylvania Certified Organic (PCO) 100% Grassfed Certification Manual, 1–10.
14. Ström G. (2012). Effect of botanically diverse pastures on the milk fatty acid profiles in New Zealand dairy cows. <http://stud.epsilon.slu.se/4257/>
15. Villeneuve M.P, Lebeuf Y., Gervais R., Tremblay G.F, Vuilleumard J.C, Fortin J., Chouinard P.Y. 2013.Milk volatile organic compounds and fatty acid profile in cows fed timothy as hay, pasture, or silage. J Dairy Sci. 96(11):7181-94.
16. Woods, V. B., & Fearon, A. M. (2009). Dietary sources of unsaturated fatty acids for animals and their transfer into meat, milk and eggs: A review. Livestock Science, 126(1–3), 1–20. <https://doi.org/10.1016/j.livsci.2009.07.002>